The Fate of Bose-Einstein Condensate in the Presence of Spin-Orbit Coupling

Qi Zhou[1] and Xiaoling Cui[2]
[1] *Department of Physics, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong*
[2] *Institute for Advanced Study, Tsinghua University, Beijing 100084, China*

(Dated: October 23, 2012)

Intensive theoretical studies have recently predicted that a Bose-Einstein condense exhibits a variety of novel properties if spin-orbit coupling is present. However, an unambiguous fact has also been pointed out that Rashba coupling destroys a condensate of non-interacting bosons even in high dimensions. Therefore, a conceptually important question arises, whether a condensate exists in the presence of interaction and a general type of spin-orbit coupling? Here we show that interaction qualitatively changes the ground state of bosons under Rashba spin-orbit coupling. Any infinitesimal repulsion forces bosons either to condense at one or two momentum states, or to form a Schrodinger-Cat state that is a superposition of infinite numbers of fragmented condensates. The Schrodinger-Cat is unstable against the anisotropy of spin-orbit coupling in systems with large numbers of particles, leading to the revival of a condensate in current experiments.

arXiv:1210.5853v1 [cond-mat.quant-gas] 22 Oct 2012

Spin-orbit coupling(SOC) is the underlying mechanism for many fundamental quantum phenomena, ranging from the atomic fine structure to the newly discovered novel properties of topological insulators[1, 2].The recent realization of synthetic SOC for neutral alkali atoms in laboratories provides physicists a new platform to study the effects of SOC in many-body systems, in which a wide range of parameters can be well controlled in experiments[3–6]. Particularly, spin-orbit coupled bosons offer physicists a unique opportunity to explore how SOC may manifest itself at the macroscopic level. As it is known as a textbook result that bosons naturally form a condensate at the ground state in three and two dimensions, intensive theoretical efforts have predicted a number of macroscopic quantum phenomena exhibited by a Bose-Einstein condensate in the presence of SOC[7–16].

On the other hand, a significant effect of SOC on non-interacting bosons has also been realized recently. It was pointed out that some types of SOC may completely destroy a non-interacting condensate at zero or any finite temperatures even in high dimensions [17–19]. To see this effect, one can start from the single-particle Hamiltonian for spin-orbit coupled bosons, which can be written as

$$\mathcal{K}=\sum_{\sigma}\int d^Dr\hat{\Psi}^{\dagger}_{\sigma}(\mathbf{r})\left(-\frac{\hbar^2\nabla^2}{2M}\right)\hat{\Psi}_{\sigma}(\mathbf{r})+H_{SOC},\qquad(1)$$

where D is the dimension, $\sigma=\uparrow,\downarrow$, $\hat{\Psi}^{\dagger}_{\sigma}(\mathbf{r})(\hat{\Psi}_{\sigma}(\mathbf{r}))$ is the creation (annihilation) operator at $\mathbf{r}$ and M is the mass. The part of SOC in the Hamiltonian is described by

$$H_{SOC}=-i\int d^Dr\hat{\Psi}^{\dagger}_{\uparrow}(\mathbf{r})(\lambda_x\partial_x-i\lambda_y\partial_y)\Psi_{\downarrow}(\mathbf{r})+c.c.,\quad(2)$$

where $\lambda_{x,y}$ is the coupling strength. For $D=3$ the low energy part of single-particle spectrum can be written as

$$\epsilon_{\mathbf{k}}=\frac{\hbar^2}{2M}(|\mathbf{k}_{\perp}|^2+k_z^2)-\sqrt{\lambda_x^2k_x^2+\lambda_yk_y^2},\qquad(3)$$

where $\mathbf{k}=(k_x,k_y,k_z)$ is the momentum and $\mathbf{k}_{\perp}=(k_x,k_y)$. For $D=2$, one can simply set $k_z=0$ in Eq.(3). Among all the configurations of SOC, Rashba coupling corresponding to $\lambda_x=\lambda_y=\lambda$ is of particular interest. In both $D=2$ and $D=3$, the kinetic energy minimum under Rashba coupling becomes a circle in the x-y plane with radius $|\mathbf{k}_{\perp}|=k_0\equiv M\lambda/\hbar^2$, which means an infinitely degeneracy of the single-particle ground state. Correspondingly, the low-energy Density of States becomes that in $(D-1)$ dimension in the absence of SOC. Therefore, a non-interacting condensate is completely destroyed at zero temperature for $D=2$[19] and at any finite temperature for $D=3$[17, 18].

Whereas the disappearance of a non-interacting condensate with Rashba coupling is rather clear, one may naturally ask whether interaction will change the fate of a condensate. To answer this question, we first note that the infinite degeneracy of single-particle ground states makes interaction effects highly non-perturbative. To determine the many-body ground state amounts to selecting the state with the lowest interaction energy from an infinitely degenerate subspace, which is known to be a challenging problem. In the literature, a mean field approach has been adopted to compare energies of a special class of states that can be described by condensate wave functions[8–14]. However, with the observation that SOC may completely destroy a condensate, one naturally concerns the validity of mean field approach. To explore the exact many-body ground state, interaction effect needs to be investigated more systematically.

We expand the interaction between two-component bosons

$$\hat{\mathcal{U}}=\Omega\int d^Dr(g_1\hat{n}^2_{\uparrow}(\mathbf{r})+g_2\hat{n}^2_{\downarrow}(\mathbf{r})+g_{12}\hat{n}_{\uparrow}(\mathbf{r})\hat{n}_{\downarrow}(\mathbf{r}))\quad(4)$$

in the basis of single-particle eigenstates, where $\hat{n}_{\sigma}(\mathbf{r})=\hat{\Psi}^{\dagger}_{\sigma}(\mathbf{r})\hat{\Psi}_{\sigma}(\mathbf{r})$ is the density operator, Ω is the volume of the system, and $g_{i=1,2}>0$, $g_{12}>0$ are the intra- and inter-spin repulsion. In this basis, the effective interac-



FIG. 1: Hartree-Fock energy $E_{HF}^{[N]}$ of N particles. Red circle represents the kinetic energy minimum, and blue ellipses represent a large number of bosons occupying the same momentum state. Blue and white dot represent a single boson and a hole in the condensate respectively. Single-particle and fragmented condensate minimize $E_{HF}^{[N]}$ for $\gamma < 1$ and $\gamma > 1$. They are separated from high energies states by an energy gap $E_G = \bar{g} + N\delta g/2$ or $E_G = \bar{g}(N/2-1)$, where $\bar{g} = (g_1+g_2+g_{12})$ and $\delta g = g_1+g_2-g_{12}$. Any two fragmented condensates with even particle numbers $N = 2n$ are coupled through a sequence of tunneling induced by n steps of scattering, as shown by the green arrows.

tion can be formulated as

$$\hat{\mathcal{U}}_L = \sum_{\phi_1,\phi_2,\phi_3,\phi_4} U_{\phi_1,\phi_2}^{\phi_3,\phi_4} \hat{L}_{\phi_1}^\dagger \hat{L}_{\phi_2}^\dagger \hat{L}_{\phi_3} \hat{L}_{\phi_4}/4, \tag{5}$$

where

$$U_{\phi_1,\phi_2}^{\phi_3,\phi_4} = g_{12} e^{i(\phi_1-\phi_4)} + g_2 + g_1 e^{i(\phi_1+\phi_2-\phi_3-\phi_4)}, \tag{6}$$

$\hat{L}_\phi^\dagger$ ($\hat{L}_\phi$) is the creation (annihilation) operator in the subspace $\mathcal{L}$ composed by all single-particle ground states, and ϕ is the polar angle on the circle $|\mathbf{k}_\perp| = k_0$[19, 20].

Using Eq.(5), a number of significant interaction effects can be revealed. First, any infinitesimal repulsion forces bosons to form a single-particle condensate or a fragmented condensate, distinct from non-interacting systems where a condensate is absent. Second, interaction inevitably builds up a superposition of fragmented condensates with even particle number, and the exact ground state becomes a Schrodinger-Cat beyond the prediction of mean field theory. Finally, Schrodinger-Cat states collapse in systems with large numbers of particle if a finite anisotropy exists in SOC, leading to the revival of a condensate in current experiments.

We start from the angular dependent $\hat{U}_L$ in Eq.(5), which drastically changes the Hartree-Fock energy of interacting systems. To see this fact, a two-body problem is an illuminating example, where Fock states can be written as either $|\phi,\theta\rangle = \hat{L}_\phi^\dagger \hat{L}_{\phi+\theta}^\dagger|0\rangle$ for $\theta \neq 0$ or $|\phi,0\rangle = \hat{L}_\phi^\dagger \hat{L}_\phi^\dagger|0\rangle/\sqrt{2}$. θ is the relative angle between the two bosons on the circle. The Hartree-Fock energy $E_{HF}^{[2]} = \langle\phi,\theta|\hat{\mathcal{U}}_L|\phi,\theta\rangle$, where the superscript $[N]$ denotes the result for a N-body problem, can be written as

$$E_{HF}^{[2]} = \begin{cases} g_{12}(1+\cos\theta)/2 + g_1 + g_2, & \theta \neq 0 \\ (g_{12}+g_1+g_2)/2. & \theta = 0 \end{cases} \tag{7}$$

Eq.(7) shows that, $|\phi,0\rangle$ and $|\phi,\pi\rangle$ minimize $E_{HF}^{[2]}$ for $\gamma < 1$ and $\gamma > 1$ respectively, where $\gamma = g_{12}/(g_1+g_2)$. States with $\theta \neq 0, \pi$ always have higher energies. For $\gamma > 1$, two bosons occupy two opposite points on the circle, i.e., fragmentation occurs. Without SOC, it is well-known that fragmentation is not possible in homogenous systems, because of the cost of Fock energy when bosons occupy different momentum states[21]. The presence of Rashba coupling suppresses the term in Fock energy that is proportional to g_{12} by a factor of $\cos\theta$. Therefore, fragmentation may occur at large enough g_{12}.

The above discussions can be directly generalized to a N-body system. For systems with even particle number, a single-particle condensate

$$|C^{[N]}\rangle_\phi = \hat{L}_\phi^{\dagger N}|0\rangle/\sqrt{N!}, \tag{8}$$

and a fragmented condensate

$$|F^{[N]}\rangle_\phi = \hat{L}_\phi^{\dagger N/2} \hat{L}_{\phi+\pi}^{\dagger N/2}|0\rangle/(N/2)! \tag{9}$$

minimize $E_{HF}^{[N]}$ for $\gamma > 1$ and $\gamma < 1$ respectively, where ϕ is arbitrary, reflecting the rotation symmetry in the momentum space for Rashba coupling. This result can be understood from that any pair of bosons prefer to occupy the same or the opposite points on the circle to save interaction energy. For odd particle number $N = 2n+1$, where n is an integer, the same conclusion holds, with the expression for the fragmented condensate simply replaced by

$$|F^{[2n+1]}\rangle_\phi = \hat{L}_\phi^{\dagger n}\hat{L}_{\phi+\pi}^{\dagger n+1}|0\rangle/\sqrt{n!(n+1)!}. \tag{10}$$

Any other Fock state in $\mathcal{L}$ costs additional Hartree-Fock energy proportional to the total particle number, as shown in Fig.1. Therefore, all bosons condense at one or two momentum states, distinct from non-interacting systems where bosons can distribute on the circle $|\mathbf{k}_\perp| = k_0$ arbitrarily.

While the significance of interaction effects has already been reflected in Hartree-Fock energy, it is worthwhile to point out that $E_{HF}^{[N]}$ only takes into account the diagonal term of interaction matrix elements. For single-particle condensates and fragmented condensate with odd particle number, such a consideration is sufficient. States $|C^{[N]}\rangle_\phi$ or $|F^{[2n+1]}\rangle_\phi$ with different values of ϕ carry different total momentum. Therefore, the minimums of Hartree-Fock energy cannot be mixed with each other by interaction due to constraint of momentum conservation. However, such a constraint is absent for fragmented condensates with even particle numbers, as all states $|F^{[2n]}\rangle_\phi$ have zero total momentum, and any two of them are inevitably coupled by a series of scattering, as demonstrated in Figure 2. This type of tunneling is known to be crucial in the presence of degenerate ground states, and produces macroscopic or mesoscopic quantum coherence in various systems[22–24]. In our case, it fundamentally changes the ground state structure when fragmentation occurs with even particle number.

We derive an effective Hamiltonian in the subspace composed by all fragmented condensates (See Appendix),

$$H_{eff}^{[2n]} = \int_0^\pi d\phi \mathcal{E}_F^{[2n]}\hat{O}_\phi^\dagger\hat{O}_\phi + \int_0^\pi d\phi d\phi' \mathcal{V}_{\phi,\phi'}\hat{O}_{\phi'}^\dagger\hat{O}_\phi(1-\delta_{\phi,\phi'}), \tag{11}$$

where $\hat{O}_\phi^\dagger = \hat{L}_\phi^{\dagger n}\hat{L}_{\phi+\pi}^{\dagger n}/n!$ is the creation operator for a fragmented condensate, and

$$\mathcal{V}_{\phi,\phi'} = (-1)^{n-1}\frac{(g_2 + g_1 e^{i2(\phi'-\phi)})^n}{(g_1+g_2+g_{12})^{n-1}}n^2. \tag{12}$$

The diagonal term $\mathcal{E}_F^{[2n]} = E_{HF}^{[2n]} + \mathcal{E}'$, where $E_{HF}^{[2n]}$ is the Hartree-Fock energy of a fragmented condensate, and $\mathcal{E}'$ is the diagonal part of the correction to the energy due to the coupling between $|F^{[2n]}\rangle_\phi$ and high energy states(See supplementary materials). As $\mathcal{E}'$ is independent on ϕ, it does not affect the structure of ground state wave function.

While $H_{eff}^{[2n]}$ is analytically solvable, the underlying physics can be revealed by considering the simplest case with $g_1 = 0$, where the system can be mapped to a one-dimension ring with an "infinite-range" constant tunneling $t = (-1)^{n-1}g_2^n n^2/(g_2+g_{12})^{n-1}$[25]. The eigenstates can be rewritten as

$$|SC_\nu^{[2n]}\rangle = \frac{1}{\sqrt{\pi}}\int_0^\pi e^{i2\nu\phi}\hat{O}_\phi^\dagger|0\rangle, \tag{13}$$

where ν is an integer, with the corresponding eigenenergy

$$E_\nu^{[2n]} = \mathcal{E}_F^{[2n]} + (-1)^{n-1}\frac{g_2^n n^2}{(g_2+g_{12})^{n-1}}(\pi\delta_{\nu,0} - 1). \tag{14}$$

As the reduced single-particle density matrix $\langle\hat{L}_\phi^\dagger\hat{L}_{\phi'}\rangle$ of $|SC_\nu^{[2n]}\rangle$ is zero, and the reduced $2n$-particle density matrix $\langle\hat{L}_\phi^{\dagger n}\hat{L}_{\phi+\pi}^{\dagger n}\hat{L}_{\phi'}^n\hat{L}_{\phi'+\pi}^n\rangle$ has a unique macroscopic eigenvalue $(n!/\pi)^2$, $|SC_\nu^{[2n]}\rangle$ is a Schrodinger-Cat state, which has distinct properties from ordinary single-particle or fragmented condensates[24]. For general cases where both g_1 and g_2 are finite, the structure of Schrodinger-Cat is similar but more elaborated(See Supplementary Materials).

It is known that Schrodinger-Cat state is fragile in systems with large numbers of particles. In our case, this very nature of Schrodinger-Cat is directly related to the fate of a condensate in the presence of SOC. Eq.(14) shows that the energy difference between the Schrodinger-Cat state and the fragmented condensate can be characterized by $\eta = 1 - E_\nu^{[2n]}/\mathcal{E}_F^{[2n]}$. For a small total particle number $N = 2n$, η takes a large value. For example, one estimates that $\eta \approx 0.61, 0.11, 0.1$ for $N = 4, 6, 8$, and $g_2/g_{12} = 0.9$, which means a large energy gained by forming a Schrodinger-Cat. While this fact strikingly changes the properties of a few-body system (See Supplementary material), one notes that η decreases exponentially with increasing particle number $N = 2n$, i.e., $\eta \sim \left(1 + \frac{g_{12}}{g_2}\right)^{-n+1}$. This is a typical feature of a high order process that requries multiple steps of two-body scattering to couple two degenerate states. When $N \to \infty$, η vanishes, which means a fragmented condensate becomes degenerate with the exact many-body ground state in the thermodynamic limit. In current experiments, $N \sim 10^5 - 10^6$, one can easily see that η is a negligible number.

The tiny η for a large value of N indicates that a big Schrodinger-Cat is unstable against external perturbation. For our discussions, the anisotropy of SOC, which always exists in practice, is naturally such a perturbation. Assuming $\lambda_x > \lambda_y$, Eq.(3) shows that kinetic energy minimums are located at two separate points $\mathbf{k} = \pm k_{x0} = \pm M\lambda_x/\hbar^2$. We define the circle $|\mathbf{k}_\perp| = k_{x0}$ as $\mathcal{L}'$, on which the expression for interaction energy is identical to that in Eq.(11), with ϕ replaced by

$\varphi = \arg\{\lambda_x k_x, \lambda_y k_y\}$[19]. The kinetic energy in $\mathcal{L}'$ is now a function of ϕ,

$$K(\phi) = \frac{\hbar^2}{2M} k_{x0}^2 - k_{x0}\lambda_x \sqrt{1 - \alpha(2-\alpha)\sin^2\phi}, \qquad (15)$$

where $\alpha \equiv (\lambda_x - \lambda_y)/\lambda_x$ characterizes the anisotropy of SOC, and leads to an offset of kinetic energy $\Delta(\alpha) = (K(\pi/2) - K(0))N = \alpha N M \lambda_x^2/\hbar^2$ in $\mathcal{L}'$.

For single-particle condensates and fragmented condensates with odd particle numbers, any infinitesimal α picks up $|C^{[N]}\rangle_{\phi=0}$ or $|F^{[N]}\rangle_{\phi=0}$ as the ground state, which favors the kinetic energy with no cost of interaction energy. For Schrodinger-Cat states, we define a characteristic anisotropy,

$$\alpha^* = \frac{\hbar^2}{2M\lambda_x^2}\left(\frac{g_2}{g_2 + g_{12}}\right)^{n-1} g_2 n, \qquad (16)$$

by setting the kinetic energy offset equal to the strength of interaction induced off-diagonal coupling, i.e., $\Delta(\alpha^*) = t$. For small anisotropy $\alpha \ll \alpha^*$, interaction energy is dominate and Schrodinger-Cat forms. For $\alpha \gg \alpha^*$, to form a superposition of fragmented condensates costs too much kinetic energy, and a single fragmented condensate $|F^{[N]}\rangle_{\phi=0}$ becomes the ground state. A more quantitative analysis is given in Supplementary Materials. Using Eq.(16), one estimates that $\alpha^* = 9\%, 4\%, 2\%$ for $N = 4, 6, 8$, $g_2/g_{12} = 0.9$ and $g_2 n\hbar^2/(2M\lambda_x^2) = 0.2$. This means a small Schrodinger-Cat is stable in a finite range of anisotropy. With increasing N, α^* decreases exponentially. One can verify that for $N \sim 10^5 - 10^6$, the typical particle number in current experiments, α^* is essentially zero. Therefore, one conclude that a Schrodinger-Cat collapses to a fragmented condensate. An additional external potential that breaks the translation invariance shall further change the fragmented condensate to a stripe condensate(See Supplemental Materials).

Whereas we have answered fundamental questions on whether, when and why a condensate exists in the presence of SOC, our work also demonstrates that the coupling between Hartree-Fock energy minimums becomes particularly important in a spin-orbit coupled few-body system. Evidence for this type of coupling has been found in a recent numerical calculation[26]. Provided that significant progresses have been made on manipulating a few atoms in current experiments[27, 28], we hope that our work will stimulate more studies on SOC induced novel ground states of mesoscopic cold atomic systems.

We thank T.L. Ho for helpful discussions. Q.Z. acknowledges start-up support from Department of Physics, The Chinese University of Hong Kong. X.C. acknowledges support from the Initiative Scientific Research Program of Tsinghua University and NSFC under Grant No. 11104158.

Appendix

We define the excited state with $n-m$ pairs of bosons at the angle ϕ and m pairs of bosons at ϕ' as $|m\rangle_{\phi,\phi'}$. Within this definition, $|F^{[2n]}\rangle_{\phi} = |0\rangle_{\phi,\phi'}$, $|F^{[2n]}\rangle_{\phi'} = |n\rangle_{\phi,\phi'}$. Interaction scatters a pair of bosons out from $|0\rangle_{\phi,\phi'}$ to the momentum states at $\mathbf{k}' = (k_0, \phi')$ and $-\mathbf{k}' = (k_0, \phi' + \pi)$. The Schrodinger equation satisfied by $|0\rangle_{\phi,\phi'}$ can be written as

$$(E_{HF}^{[2n]} - E)|0\rangle_{\phi,\phi'} + \int_0^{\pi} d\phi' \mathcal{U}_{10}|1\rangle_{\phi,\phi'} = 0, \qquad (17)$$

where $\mathcal{U}_{mn} = {}_{\phi,\phi'}\langle m|\hat{\mathcal{U}}_L|n\rangle_{\phi,\phi'}$, and E is the eigen energy. Considering the shorted path that connects $|0\rangle_{\phi,\phi'}$ and $|n\rangle_{\phi,\phi'}$ in Hilbert space as shown in Fig.(1), the Schrodinger equation satisfied by $|m\rangle_{\phi,\phi'}$ can be approximated by

$$(E_m - E)|m\rangle_{\phi,\phi'} + \sum_{l=-1}^{1} \mathcal{U}_{m+l,m}|m+l\rangle_{\phi,\phi'} = 0, \qquad (18)$$

where $E_m = {}_{\phi,\phi'}\langle m|\hat{\mathcal{U}}|m\rangle_{\phi,\phi'}$ is the Hartree-Fock energy of $|m\rangle_{\phi,\phi'}$. Eliminating $|m\rangle_{\phi,\phi'}$ from the above equations, the effective Hamiltonian in Eq.(11) that takes the standard form of a high order perturbation theory is obtained.

[1] X.-L. Qi, and S.-C. Zhang, Rev. Mod. Phys. 83, 1057 (2011)

[2] M. Z. Hasan, and C. L. Kane, Rev. Mod. Phys. 82, 3045 (2010)

[3] Y.-J. Lin, K. Jiménez-García, and I. B. Spielman, Nature 471, 83-86 (2011)

[4] L. W. Cheuk, A.T. Sommer, Z. Hadzibabic, T. Yefsah, W. S. Bakr, and M.W. Zwierlein, Phys. Rev. Lett. 109, 095302 (2012)

[5] J. Y. Zhang, S.C. Ji, Z. Chen, L. Zhang, Z.D. Du, B. Yan, G.S. Pan, B. Zhao, Y.J. Deng, H. Zhai, S. Chen, and J.W. Pan, Phys. Rev. Lett. 109, 115301 (2012)

[6] P. Wang, Z.Q. Yu, Z. Fu, J. Miao, L. Huang, S. Chai, H. Zhai, and J. Zhang, Phys. Rev. Lett. 109, 095301 (2012)

[7] T.-L. Ho, and S. Zhang, Phys. Rev. Lett. 107, 150403, (2011).

[8] C. Wang, C. Gao, C.M.Jian, and H. Zhai, Phys. Rev. Lett. 105, 160403 (2010).

[9] T. D. Stanescu, B. Anderson, and V. Galitski, Phys. Rev. A 78, 023616 (2008)

[10] C.Wu, I. Mondragon-Shem, and X-F. Zhou, Chin. Phys. Lett. 28, 097102 (2011)

[11] H. Hu, B. Ramachandhran, H. Pu, and X-J. Liu, Phys. Rev. Lett. 108, 010402(2012)

[12] S. Sinha, R. Nath, and L. Santos, Phys. Rev. Lett. 107, 270401 (2011)

[13] T. Ozawa, and G. Baym, Phys. Rev. A 85, 013612 (2012)

[14] R. Barnett, S. Powell, T. Grass, M Lewenstein, and S. Das Sarma, Phys. Rev. A 85 023615 (2012)

[15] Y. Li, L.P. Pitaevskii, and S. Stringari, Phys. Rev. Lett. 108, 225301 (2012)

[16] Z. F. Xu, Y. Kawaguchi, L. You, and M. Ueda, arXiv:1203.2005 (2012)

[17] H. Hu, and X.-J. Liu, Phys. Rev. A 85, 013619 (2012)
[18] T. Ozawa, and G. Baym, Phys. Rev. Lett. 109, 025301 (2012)
[19] X. Cui, and Q. Zhou, arXiv:1206.5918 (2012)
[20] T. Ozawa, and G. Baym, Phys. Rev. A 84, 043622 (2011)
[21] P. Nozieres, Some Comments on Bose-Einstein Condensation. Bose-Einstein Condensation, edited by A. Griffin, D.W. Snoke, and S. Stringari (Cambridge University Press, Cambridge, 1995)
[22] A.J. Leggett, Quantum Tunneling in Condensed Media, edited by Y. Kagan and A.J. Leggett, (Elsevier Science 1992)
[23] W. Wernsdorfer, and R. Sessoli, Science 284, 133 (1999)
[24] T.L. Ho, and C. V. Ciobanu, J. Low Temp. Phys. 125257 (2004).
[25] P. van Dongen, and D. Vollhardt, Phys. Rev. B 40, 7252 (1989)
[26] B. Ramachandhran, B. Opanchuk, X-J. Liu, H. Pu, P. D. Drummond, and H. Hu, Phys. Rev. A 85, 023606 (2012)
[27] F. Serwane, G. Zrn, T. Lompe, T. B. Ottenstein, A. N. Wenz, and S. Jochim, Science 332, 336 (2011)
[28] W. Bakr, A. Peng, M. Tai, R. Ma, J. Simon, J. Gillen, S. Foelling, L. Pollet, and M. Greiner, Science 329, 547 (2010)

SUPPLEMENTARY MATERIAL

In this supplementary material, we present the results on the diagonal term of Eq.(4) in the main text, the general solution for the effective Hamiltonian H_{eff}, small N systems, effects of anisotropy of spin-orbit coupling, and the evolution from a fragmented condensate to a stripe condensate.

Diagonal term of Eq.(4)

In Eq.(4) of the main text, the Hartree-Fock energy can be written as

$$E_{HF}^{[2n]} = \frac{1}{4}(6g_1 + 6g_2 + 2g_{12})n^2 - \frac{1}{2}(g_1 + g_2 + g_{12})n, \quad (19)$$

For $\mathcal{E}'$, to the second order approximation for the diagonal term, we consider the lowest excited states with only one pair of atoms scattered out from the fragmented condensate and obtain

$$\mathcal{E}' = \int_0^\pi d\phi' |\langle Ex|\hat{\mathcal{U}}_L|F^{[2n]}\rangle|^2/E_G, \quad (20)$$

where $|Ex\rangle = \hat{L}_\phi^{\dagger n-1}\hat{L}_{\phi+\pi}^{\dagger n-1}\hat{L}_{\phi'}^\dagger\hat{L}_{\phi'+\pi}^\dagger|0\rangle/(n-1)!$ and $E_G = \langle Ex|\hat{\mathcal{U}}_L|Ex\rangle - E_{HF}^{[2n]} = (g_1 + g_2 + g_{12})(n-1)$. To be explicit, $\mathcal{E}' = \pi\frac{n^2}{n-1}\frac{g_1^2+g_2^2}{g_1+g_2+g_{12}}$.

Solution of H_{eff}

For general cases of $g_1 \neq 0, g_2 \neq 0$, $\mathcal{V}_{\phi,\phi'}$ can be written as

$$\mathcal{V}_{\phi,\phi'} = \frac{(-1)^{n-1}n^2}{(g_1 + g_2 + g_{12})^n}\sum_{\kappa=0}^{n}\mathcal{C}_n^\kappa g_1^\kappa g_2^{n-\kappa}e^{2i\kappa(\phi'-\phi)}. \quad (21)$$



FIG. 2: Many-body ground states under Rashba coupling. single-particle condensate occupies the region with small γ. For large γ, the ground state is a fragmented condensate at odd particle number, $N = 4l + 1$ or $N = 4l + 3$, where l is an integer. For even particle numbers, the ground state is a Schrodinger-Cat state with infinite degeneracy for (I) with $N = 4l + 2$ and no degeneracy for (II) with $N = 4l$. The dashed red curve represents the critical value γ_c for the transition between single-particle condensate and other three states.

It is straightforward to verify that $|SC_\nu^{[2n]}\rangle$ are the eigen states with the eigen energies

$$E_\nu^{[2n]} = \begin{cases} \mathcal{E}_F^{[2n]} + \frac{(-1)^{n-1}n^2}{(g_1+g_2+g_{12})^{n-1}}(\pi\mathcal{C}_n^\nu g_1^\nu g_2^{n-\nu} - (g_1 + g_2)^n) \\ \quad 0 \leq \nu \leq n \\ \\ \mathcal{E}_F^{[2n]} - (-1)^{n-1}\frac{(g_1+g_2)^n n^2}{(g_2+g_{12})^{n-1}} \\ \quad \nu < 0, \nu > n \end{cases} \quad (22)$$

where $\mathcal{C}_n^\nu = \frac{n!}{\nu!(n-\nu)!}$. For $N = 2n = 4l + 2$, $(-1)^{n-1} > 0$, all states with $\nu > n$ are degenerate and have lower energy than those with $\nu \leq n$. This means the ground states are infinitely degenerate. For $N = 2n = 4l$, $(-1)^{n-1} < 0$, the ground state is unique, which is determined by the ratio of g_1/g_2 so that $\mathcal{C}_n^\nu g_1^\nu g_2^{n-\nu}$ is maximized. For example, for $g_1/g_2 = 0, 1, \infty$, the winding number for the ground state $\nu_0 = 0, n/2, n$ respectively.

Small N systems

The dependence of the ground state on the total particle number and interaction is summarized in Fig.(2). $\gamma_c(N)$ is the boundary between different phases, which is determined by comparing the energy of the single-particle condensate $|C^{[2n]}\rangle$ with other states. For instance, by solving $E_C^{[2n]} = E_\nu^{[2n]}$, one obtains the value of $\gamma_c(N)$ that separates $|C^{[2n]}\rangle$ and the Schrodinger-Cat $|SC^{[2n]}\rangle$.

At small particle numbers, the derivation of $\gamma_c(N)$

from the mean field value $\gamma_c^0 = 1$ is significant. For $N = 2n = 2$, this two-body problem can be solved exactly. Under this situation, $\mathcal{E}'$ is absent, and $\mathcal{E}_F^{[2]} = E_{HF}^{[2]} = g_1 + g_2$. By setting $n = 1$ in Eq.(22), it can be verified easily that for any $\nu \neq 0, 1$, the eigen energy is exactly zero. Comparing with the energy of any state with a finite total momentum, we see that the ground state is always a small Schrodinger-Cat state, regardless of the ratio between g_{12} and $g_1 + g_2$.

In spite of the repulsive interaction, we see that the ground state energy of two bosons for Rashba coupling is exactly zero. The underlying physics is that the scattering of a pair of bosons with zero momentum on the circle cancels the positive Hatree-Fock energy. It is interesting to note that, this result is consistent with the conclusion of a completely different approach based on renormalization of two-body interaction, which discovered that the effective interaction between two particles on the opposite points of the circle is zero[20].

It is also useful to study how $\gamma_c(N)$ approaches the mean field value at large N. Clearly, the contribution from $\mathcal{E}'$ to the ground state energy is much larger than that from off-diagonal term in Eq.(4) of the main text. If one ignores $\mathcal{E}'$, $E_C^{[2n]} = \mathcal{E}_F^{[2n]}$ leads to the mean-field result $\gamma_c^0 = 1$. Taking into account $\mathcal{E}'$, the value of γ_c is modified. In large N limit, we find

$$\gamma_c(2n) \approx \gamma_c^0 - \frac{\pi}{n}\frac{g_1^2 + g_2^2}{(g_1 + g_2)^2}. \tag{23}$$

Similar scaling of $\gamma_c(N) - \gamma_c^0 \sim 1/N$ holds for odd particle numbers.

Effect of anisotropy of SOC

The kinetic energy offset in $\mathcal{L}'$ as shown in Eq. (8) of the main text leads to an additional term in the Hamiltonian for anisotropic SOC,

$$\hat{\mathcal{W}} = \int_0^\pi d\phi K(\phi) \hat{L}_\phi^\dagger \hat{L}_\phi. \tag{24}$$

The matrix elements of $\hat{\mathcal{W}}$ between $|SC_\nu^{[2n]}\rangle$ can be calculated straightforwardly,

$$\mathcal{W}_{\nu,\nu'} = \frac{N}{\pi}\int_0^\pi d\phi e^{2i(\nu-\nu')\phi} K(\phi) \tag{25}$$

Since $K(\phi + \pi) = K(\phi)$, $\mathcal{W}_{\nu,\nu'}$ is finite only when $\nu - \nu'$ is an integer. For Rashba coupling, $K(\phi)$ is a constant and therefore $\mathcal{W}_{\nu,\nu'} = 0$. With increasing $\delta K = K(\pi/2) - K(0)$ or equivalently the anisotropy of SOC α, the amplitude of $\mathcal{W}_{\nu,\nu'}$ increases.

Tthe general expression of the ground state wave function can be written as

$$|SC^{[2n]}\rangle_\alpha = \int_0^\pi d\phi C_\phi \hat{O}_\phi^\dagger |0\rangle. \tag{26}$$

For isotropic SOC and $N = 2n = 4l$, $C_\phi = e^{2i\nu_0\phi}/\sqrt{\pi}$, and $|C_\phi|^2 = 1/\pi$ is a constant, where $\nu_0 \leq n$ as discussed before. In the presence of a small anisotropy, $\hat{\mathcal{W}}$ is finite, one may take into account only the mixing between $|SC_{\nu_0}^{[2n]}\rangle$ and $|SC_{\nu_0\pm1}^{[2n]}\rangle$, which leads to a small variance of $|C_\phi|^2$. With increasing α, more states with different ν are mixed into the ground state. As $\mathcal{W}_{\nu,\nu'} \approx N\delta K$, we see for large enough $N\delta K \gg E_{\nu>n} - E_{\nu_0}$, all states $|SC_\nu^{[2n]}\rangle$ are mixed. An equal superposition of them leads to $C_\phi = \delta_{\phi,0}/\sqrt{\pi}$ and the ground state becomes a fragmented condensate.

For $N = 2n = 4l + 2$, a subtle difference emergies due to the infinitely degenerate ground states under Rashba coupling. In the presence of any infinitesimal $\hat{\mathcal{W}}$, a superposition of these degenerate state $|SC_{\nu>n}^{[2n]}\rangle$ may fulfill the goal of minimizing both the kinetic energy and interaction energy. For instance, in the simplest case where $g_1 = 0$, the ground state becomes

$$|SC^{[2n]}\rangle_{\alpha\to0} = \sum_{\nu\neq0} |SC_\nu^{[2n]}\rangle = \int_0^\pi C_\phi \hat{O}_\phi^\dagger |0\rangle, \tag{27}$$

where $C_\phi = (\pi\delta_{\phi,0} - 1)/\sqrt{\pi^2 - \pi}$. With a finite α, $|SC_0^{[2n]}\rangle$ will be mixed to the ground state. For large enough $N\delta K \gg E_{\nu>n} - E_{\nu_0}$, similar to the case of $N = 2n = 4l$, all states $|SC_\nu^{[2n]}\rangle$ are mixed and a fragmented condensate becomes the ground state.

Fragmented condensate to a stripe condensate

The fragmened condensate is a Fock state in the momentum space, with half particles occupying the state $\mathbf{k_0}$ and the other half occupying $-\mathbf{k_0}$. In homogenous systems, these two states can not be mixed due to momentum conservation. If an external potential exists and translational invariance is broken, an additional term exists in the Hamiltonian $H' = t'\hat{L}_{\mathbf{k_0}}^\dagger \hat{L}_{-\mathbf{k_0}} + c.c$, where t' is proportional to the strength of the potential. Therefore, it becomes a double-well problem in momentum space. It is well known that the tunneling between the two "wells" is enhanced with increasing particle number N due to the bosonic enhancement effect, i.e., $H' \sim t'N$. In the thermodynamic limit, any infinitesimal t' leads to the formation of a coherent state $\sim (\hat{L}_{\mathbf{k_0}}^\dagger + \hat{L}_{-\mathbf{k_0}}^\dagger)^N |0\rangle$, so called stripe condensate in the literature.